

WEISMAN CHILDREN'S REHABILITATION HOSPITAL
Pharmacy Manual

SUBJECT: House Stock Medications

No: 216

EFFECTIVE DATE: 12/97

Page: 1 OF 1

REVIEWED/REVISED: 4/03, 4/04, 6/05, 6/06, 6/07, 6/08, 3/09, 8/11, 6/12, 6/13,
6/14, 1/15, 10/17, 8/20, 9/23

PURPOSE:

To delineate stock medications, ordering procedure, and administration guidelines.

PROCEDURE:

1. Only non-legend drugs are maintained as stock. They must be permanently labeled to include the drug name, manufacturer, lot number, expiration date and cautionary or ancillary labeling.
2. Stock medications may be administered only on the order of a Licensed Physician. All rules and regulations pertaining to the administration of medications are also applicable to stock medications.
3. Stock medications are ordered from pharmacy via Pharmacy Order Sheet, indicating by order they are floor stock.

House Stock/Back Up Meds

Appendix A: List of House Stock Meds.

Medical Director

Provider Pharmacy

Nurse Executive

Consultant Pharmacist

Appendix A

**WEISMAN CHILDREN'S REHAB HOSP****BACK UP BOX CONTENTS**

ANTIBIOTICS	CAUTIONARY	PAR LEVEL
Amoxicillin (Amoxil) 250 mg/5 mL Susp	May be taken with food.	1 x 100 mL
Amoxicillin (Amoxil) 250 mg Cap	May be taken with food.	5
Amoxicillin/Clavulanate (Augmentin) 250 mg/5 mL Susp	May give without regard to meals.	1 x 100 mL
Amoxicillin/Clavulanate (Augmentin) 250 mg Tab	May give without regard to meals.	5
Amoxicillin/Clavulanate (Augmentin) 400 mg Tab	May give without regard to meals.	5
Cefuroxime (Ceftin) 250 mg Tab	May give with or without food.	5
Cephalexin (Keflex) 250 mg Cap	May give without regard to meals.	5
Cephalexin (Keflex) 250 mg/5 mL Susp	May give without regard to meals.	1 x 100 mL
Ciprofloxacin (Cipro) 250 mg Tab	Administer at least 2 hours before or 6 hours after magnesium/aluminum antacids and other products containing calcium, iron, or zinc.	5
Clarithromycin (Biaxin) 250 mg Tab	May give with or without food. Administer every 12 hours rather than twice daily to avoid peak and trough variation.	5
Clindamycin (Cleocin) 75 mg/5 mL Soln	May take with food.	1 x 100 mL
Metronidazole (Flagyl) 250 mg Tab	May give with food to minimize stomach upset.	5
Sulfamethoxazole-TMP (Bactrim) Susp	Take with at least 8oz water.	1 x 473 mL
Sulfamethoxazole-TMP (Bactrim) 400/80 mg Tab	Take with at least 8oz water.	5
Sulfamethoxazole-TMP (Bactrim DS) 800/160 mg Tab	Take with at least 8oz water.	5
Tobramycin 80 mg/2 mL Inj Vial	Stable at room temperature as clear, colorless solution and as dry powder.	2
NON-ANTIBIOTICS	CAUTIONARY	PAR LEVEL
Acetaminophen (Tylenol) 160 mg/5 mL UD Cup		10 x 5mL
Acetaminophen (Tylenol) 120 mg Supp		12
Acetaminophen (Tylenol) 325 mg Supp		12
Acetaminophen (Tylenol) 325 mg Tab		10
Aspirin 81mg Chew Tab		10
Bisacodyl (Dulcolax) 10mg Supp		12
Diphenhydramine (Benadryl) 25 mg Cap	May cause drowsiness. Avoid alcohol.	5
Diphenhydramine (Benadryl) 50 mg/mL Inj Vial	May cause drowsiness. Avoid alcohol.	2 x 1mL
Docusate (Colace) 50 mg/5 mL UD Cup		10 x 10mL
Docusate (Colace) 50 mg Cap		10
Epipen JR 0.15 mg Inj Syr 2-Pak		1 box
Fleet Enema (Adult)		2 x 113 mL
Fleet Pedia-Lax Enema (Pediatric)		2 x 66 mL
Enoxaparin (Lovenox) 30 mg/0.3 mL Inj Syr	Administer by deep SubQ injection alternating between the left or right anterolateral and left or right posterolateral abdominal wall. Do not mix with other infusions or injections. In order to minimize bruising, do not rub injection site. To avoid loss of drug from the 30 mg and 40 mg prefilled syringes, do not expel the air bubble from the syringe prior to injection.	2
Enoxaparin (Lovenox) 60 mg/0.6 mL Inj Syr		2
Enoxaparin (Lovenox) 100 mg/mL Inj Vial		1 x 3 mL

WEISMAN CHILDREN'S REHAB HOSP**BACK UP BOX CONTENTS**

NON-ANTIBIOTICS	CAUTIONARY	PAR LEVEL
Furosemide (Lasix) 10 mg/mL Soln	May be administered with food or milk if GI distress occurs.	1 x 60 mL
Glucose 4 gm Chew Tab		1 x 10
Glucose (Insta-Glucose) Gel		5 x 31 gm
Glycerin Pediatric Supp		12
Glycerin Adult Supp		12
Guaifenesin (Robitussin) 100 mg/5 mL UD Cup		10 x 5 mL
Ibuprofen (Motrin) 100 mg/5 mL UD Cup	Administer with food or milk.	10 x 5 mL
Ibuprofen (Motrin) 200 mg Tab	Administer with food or milk.	10
Mag-Al Plus UD Cup		10 x 30mL
Milk of Magnesia UD Cup		10 x 30mL
Prednisone 5 mg/5 mL Soln	Take with food or milk.	1 x 120 mL
Prednisone 1 mg Tab	Take with food or milk.	10
Prednisone 10 mg Tab	Take with food or milk.	10
Protamine 50 mg/5 mL Inj Vial		2 x 5mL
Silver Nitrate App		1 x 100
Sodium Chloride 0.9% Flush Syr		5 x 10 mL
Sterile Water for Injection Vial		5
Warfarin (Coumadin) 2 mg Tab	Do not take aspirin or aspirin products.	2
Warfarin (Coumadin) 2.5 mg Tab	Do not take aspirin or aspirin products.	2
Warfarin (Coumadin) 3 mg Tab	Do not take aspirin or aspirin products.	2
Warfarin (Coumadin) 4 mg Tab	Do not take aspirin or aspirin products.	2
Warfarin (Coumadin) 5 mg Tab	Do not take aspirin or aspirin products.	2
INHALED MEDICATIONS	CAUTIONARY	PAR LEVEL
Albuterol 0.083% (2.5 mg/ 3mL) Inh Soln	Do not use if solution changes color or becomes cloudy. Products packaged in foil should be used within 1 week (or according to the manufacturer's recommendations) if removed from foil pouch.	1 box
Levalbuterol (Xopenex) 0.63mg Inh Soln	Vials should be used within 2 weeks after opening protective pouch. Use within 1 week and protect from light if removed from pouch.	1 box
Levalbuterol (Xopenex) 1.25mg Inh Soln	Vials of concentrated solution should be used immediately after removing from protective pouch.	1 box
ProAir HFA	Shake well before use; prime prior to first use, and whenever inhaler has not been used for >2 weeks or when it has been dropped, by releasing 3 to 4 test sprays into the air (away from face).	1 inhaler